


# ARE YOU AND YOUR FAMILY UP TO DATE?

## COVID-19 Vaccine Schedule for age 6 Months and Up


for People who **ARE** Moderately or Severely Immunocompromised (weakened immune system)




### 6 months - 4 years




OR




### 5 years



OR



### 6 - 11 years



**For Ages 12+,  
Check the back!**



Having a weakened immune system means you are at **greater risk** of getting sick with COVID-19 so it is especially important to keep up to date with vaccines to stay protected.


\* For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.



## 12 years and older




### Moderna or Pfizer-BioNTech



OR

### Novavax



## 18 years and older who previously received Janssen primary series dose ‡



If you are moderately or severely immunocompromised or severely allergic to COVID-19 vaccines, talk to your healthcare provider about EVUSHELD™, which is a medicine that can help prevent you from getting COVID-19.

\* For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.

† A monovalent Novavax booster dose may be used in limited situations in persons age 18+ who completed a primary series using any COVID-19 vaccines, have not received a previous booster dose(s) and are unable or unwilling to receive an mRNA vaccine. It is given at least 6 months after completion of primary series.

‡ Janssen COVID-19 Vaccine should only be used in certain limited situations.

See <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a>

**NJ COVID-19  
Call Center:  
1-855-568-0545  
M-F 10am-6pm;  
Saturday 10am-4pm**

**Or scan for the nearest site!**

